

# Stray Creek Wildlife Effects

## Relevant Laws, Regulations, and Policy and Compliance

### Regulatory Framework

#### Clearwater National Forest Land and Resource Management Plan

The 1987 Forest Plan documents goals, objectives, and standards, for managing Forest wildlife species and habitats. Forest Plan goals (page II-2-2) provide for viable populations of all indigenous wildlife species, maintain and improve habitat for big-game, limit motorized use on selected big-game range, and manage habitat to contribute to the recovery of threatened and endangered species on the Forest. Objectives (FP page II-2-5) focus on maintenance and improvement on big-game ranges. Forest Plan standards are on page II-23-24.

**Table 1. Stray Creek project consistency with Clearwater Forest Plan standards**

Standard number	Standard Summary	Project Compliance Achieved By
Forest Wide Standards		
Wildlife and Fish		
a.	Provide the proper mix of hiding and thermal cover, forage, and protection from harassment during critical periods on big-game summer range (primarily elk) in accordance with criteria contained in the "Guidelines for Evaluating and Managing Summer Elk Habitat in Northern Idaho."	The Project would increase opening for big-game summer range by 11% in the affected area. Elk analysis details are available in the project record. The affected elk analysis areas meets the Forest Plan Standards of maintaining elk habitat (see Elk effects).
c.	Provide habitat for snag-dependent indicator species (pileated woodpecker and goshawk) in accordance with guidelines provided in Appendix H.	Old growth and snag retention is maintained in project area as described in the description of the proposed action. See Forest Vegetation analysis for meeting old growth standards.
f.	Provide an adequate amount of habitat to support the Clearwater Forest's assigned goal of ten endangered gray wolves as based on recommendations from the Northern Rocky Mountain Recovery Team.	Habitat is maintained or altered to provide increased big game forage over all seasons; increasing the prey base for wolves.
g.	Cooperate with future recovery efforts on behalf of the gray wolf, bald eagle, and grizzly bear.	Habitat is managed for increased forage for big game. This would provide an increased prey base for the wolf and bear. Grizzly bear is not considered as a species with occupied habitat on the Forest. The project area is not considered as nesting habitat for the bald eagle.
j.	Cooperate with Idaho Fish and Game, Indian tribes, and other agencies in the management of wildlife and fish habitat.	Idaho Fish & Game, other local agencies, and Nez Perce Tribe have been notified of project activities through scoping and informal consultation.
Timber		
j.	Manage tree openings created by	The Project would increase opening for

Standard number	Standard Summary	Project Compliance Achieved By
	<p>even-age timber harvest as follows:  (3) Duration of openings - consider an opening no longer an opening when the density and height of the vegetation and watershed conditions meet the resource management objectives of the area.  Big-Game Summer Range/Timber - In proposed EI and E3 Management Areas. the minimum standard is to provide 25 percent elk habitat potential. New openings (regeneration cuts) can be planned adjacent to former openings as long as the former opening is certified as stocked and the area meets a minimum of 25 percent elk habitat potential after implementation of the proposed activity.  The ID Team must assure that unit design optimizes wildlife objectives, both short-and long-term, within the overall objectives of the management area. Other resource requirements and objectives such as visual, watershed, silvicultural, etc., also must be met as applicable. The dispersal of timber size class objectives in the Regional Guide must be met.</p>	<p>big-game summer range by 11% in the affected area. Elk analysis details are available in the project record. The affected elk analysis area meets the Forest Plan Standards of maintaining elk habitat (see Elk effects).</p>
Management Area E1		
Wildlife and Fish b.	<p>Manage for a minimum of 25 percent maximum elk potential habitat effectiveness. During Plan implementation and further analysis, determine whether remaining areas of EI have potential for providing elk habitat. When analysis shows elk potential is limited by factors other than National Forest management, determinations may be made not to manage for elk. When habitat conditions warrant, managers are urged to exceed the 25 percent habitat standard. See Forestwide General Standards, in Chapter II</p>	<p>The proposed action would increase opening for big-game summer range by 11% in the affected area. Elk analysis details are available in the project record. The affected elk analysis area meets the Forest Plan Standards of maintaining elk habitat (see Elk effects).</p>
Timber c.	<p>Identify and maintain suitable old-growth stands and replacement habitats for snag and old-growth dependent wildlife species in accordance with criteria in Appendix H.</p>	<p>Old growth and snag retention is maintained in project area as described in the description of the proposed action. See Forest Vegetation analysis for meeting old growth standards.</p>

## Federal Law

### *Endangered Species Act*

No habitat or occurrence of threatened or endangered species is present in the Stray Creek project area.

### *Migratory Bird Treaty Act and Executive Order 13186*

The Migratory Bird Treaty Act (MBTA) prohibits taking of migratory birds, their parts, nests, eggs, and nestlings. Executive Order 13186, signed January 10, 2001, directs federal agencies to protect migratory birds by integrating bird conservation principals, measures, and practices into agency activities and to avoid or minimize, to the extent practicable, adverse impacts on migratory bird resources when conducting agency actions. Additional direction comes from the MOU between the Forest Service and USFWS, signed December 2008. The purpose of this MOU is to strengthen migratory bird conservation through enhanced collaboration between the Forest Service and USFWS, in coordination with State, tribal, and local governments.

The proposed action is in compliance with the MBTA and Executive Order 13186, which authorizes activities including habitat protection, restoration, enhancement, necessary modification, and implementation of actions that benefit priority migratory bird species (Memorandum of Understanding Between USDA Forest Service and USDI Fish & Wildlife Service – 01-MU-11130117-028), described further below).

In late 2008, a Memorandum of Understanding between the USDA Forest Service and the US Fish and Wildlife Service to Promote the Conservation of Migratory Birds was signed. The intent of the MOU is to strengthen migratory bird conservation through enhanced collaboration and cooperation between the Forest Service and the Fish and Wildlife Service as well as other federal, state, tribal and local governments. Within the National Forests, conservation of migratory birds focuses on providing a diversity of habitat conditions at multiple spatial scales and ensuring that bird conservation is addressed when planning for land management activities.

Project activities have the potential to affect migratory birds by altering habitat and displacing birds through disturbance. In areas where activities are ongoing, breeding birds may avoid or abandon habitats to avoid human activities and disturbance. Activities would be limited in time and spatial extent, so effects would be temporary and on a small scale. Proposed activities would not affect migratory birds at the planning unit scale. The project will maintain a mosaic of vegetation types and age classes to provide for a diversity of species, meets the requirements for snags and old growth (Appendix H of Forest Plan). The proposed action was designed to protect or enhance priority habitats for landbird species, including neotropical migratory species. Design criteria for project activities cover potential disturbances to birds, and allow for mitigations of the project if necessary.

### *National Forest Management Act*

This act requires the Forest Service to provide for diversity of plant and animal communities based on the suitability and capability of the specific land area in order to meet overall multiple-use objectives (16 USC 1604(g)(3)(B)). The Forest Service's focus for meeting the requirement of NFMA and implementing its regulations depends on assessing habitat to provide for diversity of species. The proposed action would be consistent with NFMA direction for diversity of animal communities. Although the proposed action analyzed in the Project may impact individual animals, the Project would not affect the viability of any species across its range.

**Sensitive Species:** Sensitive wildlife species are those that show evidence of a current or predicted downward trend in population numbers or habitat suitability that would substantially reduce species distribution. Federal laws and direction applicable to sensitive species (SS) include the NFMA and FSM 2670. The Forest is required to determine the potential effect of proposed activities on SS and to prepare biological evaluations. The Forest Service is bound by federal statutes (ESA, NFMA), regulations, and agency policy (FSM 2670) to conserve biological diversity on NFS lands and assure sensitive species populations do not decline or trend toward listing under the ESA. This document fulfills the requirements of the biological evaluation for sensitive species. The Proposed Actions would not affect sensitive species viability on federal lands, nor would it cause sensitive species to become federally listed as threatened or endangered.

**Species Viability:** The proposed action, in combination with and within the context of past, present, and reasonably foreseeable future management actions in the analysis area, would not affect population viability or distribution of native and desired nonnative vertebrate species on the Forest. At the Forest-wide scale, this Project would not disturb, agitate or bother populations to a degree that causes, or is likely to cause, a measurable decrease in productivity by substantially interfering with normal breeding, feeding, or sheltering behavior.

This analysis incorporates the effects on terrestrial sensitive species and fulfills the requirements of the required Biological Evaluation, per direction pertaining to the FSM and streamlining process (USDA Forest Service 1995). The streamlined process for doing biological evaluations for sensitive species focuses on the following two areas:

- Incorporating the Effects on Sensitive Species into the NEPA Document
- Summarizes the Conclusions of Effects of the Biological Evaluations for Sensitive Species

## Data Sources

Modeled vegetation layers (from GIS applications) were used for interpretation of species habitats or potential habitat for the animals' life stages. Data related to vegetative features as potential wildlife species habitat was queried from the vegetation model V-Map, and other data on riparian, road, fire and harvest layers.

*“VMap is a remote sensed product which uses a combination of satellite imagery and airborne acquired imagery. The image data (i.e., pixels) are put through a process of aggregation to derive spatially cohesive units (i.e., polygons). A small portion of these polygons are then sampled through aerial photo interpretation and field data collection to determine their composition and through spatial statistics, unsampled polygons are given labels based on an analysis of the sampled polygons. Draft map products are then field verified and appropriate changes are made in the labeling algorithms. Final results are then used to populate the VMap base-level feature class. A variety of post-processing algorithms are then used to create the mid-level feature classes of the VMap database.”* (Brown and Barber 2012). A follow up accuracy assessment of the following variables used in V-Map was estimated at 74% accuracy; produced by Region 1 (Brown 2015).

The most recent monitoring report is located in the project record provides information on species includes modelled habitat runs, references to other surveys or reports conducted by state, federal or private interests and monitoring surveys (document 23-083).

## Species Dropped from Detailed Analysis

The following species were dropped from detailed analysis as suitable habitat is not present in or near activity areas, or the project would not affect individuals or their habitats: bald eagle, belted kingfisher, black-backed woodpecker, Canada lynx, Coeur d' Alene salamander, flammulated owl, fringed myotis, grizzly bear, harlequin duck, pygmy nuthatch, ringneck snake, Townsend's big-eared bat, and wolverine.

## Affected Environment

### Fisher

Fishers are associated with mature coniferous forests and specific structural elements; particularly large trees and coarse woody debris (Samson, 2006b; Ruggiero et al. 1994). They inhabit mesic, coniferous forest between 3,500–6,000 feet elevation, although habitat preference changes with season, age, and sex (Badry, 2004; NatureServe, 2019). Some researchers found that fishers did not select dry forest types with large representation of ponderosa or lodgepole pines (Schwartz et al. 2013, Olson et al. 2014). Many authors mention that fisher avoids open areas. Weir and Corbould, 2010; Schwartz et al. 2013; Sauder 2014; and Sauder and Rachlow, 2014). Examples of such open areas include, grassy openings, meadows, and recently logged or severely burned areas within the past 15 years.

Fishers have a preference for structurally complex areas with multiple canopy layers, including understory shrubs and large amounts of woody debris. Ruggiero et al. (1994) concluded that riparian zones, high elevation old growth grand fir, and subalpine fir stands are important habitat components for fisher. Another set of researchers found that high elevation forested stands of lodgepole pine and sub-alpine fir were not considered as fisher habitat (Olsen et al. 2014). Fishers use "many different habitats for hunting as long as these areas provide overhead cover at either the stand or patch scales" (Weir and Harestead, 2003). Sufficient overhead cover in foraging habitat may be provided by either tree or shrub cover. Although fisher home ranges are consistently characterized by moderate to high proportions of mid- and late seral forests, there are few overarching patterns of selection for particular seral conditions or species compositions (Sauder and Rachlow, 2014). Raley et al. (2012) hypothesized that when fishers select home ranges, they benefit from including a diverse array of available forest conditions by increasing access to a greater diversity and abundance of prey species while still attaining habitat features important for reproduction and thermoregulation. Sauder and Rachlow's (2014) results are consistent with this contention and estimated an average male home range at around 24,300 acres, and a female home range of about 12,200 acres.

For this analysis, potentially suitable habitat was determined using a model (Sauder, 2014) that combines three models of fisher habitat including: a climate model (Olson et al. 2014), a landscape-scale model (Sauder and Rachlow, 2014), and a home range scale model (Sauder and Rachlow, 2015). The Sauder (2014) model identifies relative probability of fisher occurrence and its continuity across the landscape. GIS modelled runs determined open areas, probable/general habitat and mature habitat. Open areas consisted of tree canopy cover at 10% or less: and were considered as habitats the fisher usually avoids (Sauder, 2014). Open areas contribute to fragmentation of the desired canopy cover for fisher. Fragmented open habitat consists of isolated openings that are surrounded by probable fisher habitat. When these open areas reach a

cumulative 5% or greater representation in the project area, they contribute to fragmentation of fisher habitat that is less than optimal for the predator (Sauder 2014, Sauder and Rachlow 2014).

Stands classified as having a mature, mesic-mixed conifer forest (determined by the species with the greatest abundance of canopy cover, basal area, or trees per acre) were selected and intersected with “probable habitat” by the Sauder (2014) model. Selection criteria for mature forests were those areas greater than 10 inches DBH.

In summary, the most current science for the Nez Perce-Clearwater National Forest recommends landscapes that have greater than 50% mature forest arranged in contiguous, complex shapes with few isolated patches, and open areas comprising less than 5% of the area appear to constitute a forest pattern occupied by fishers (Sauder 2014, Sauder and Rachlow 2014).

Forest direction for analysis of fisher territories is to use a subwatershed (Hydrologic Unit Compartment (HUC) at the size of 12 (10,000-40,000 acres)). The Middle Lolo subwatershed (HUC12) used for analyzing a fisher territory is over 29,000 acres in size. The amount of modelled fisher habitat within the analysis area was calculated at 3,916 acres of mature habitat and 14,137 acres of general fisher habitat. Over 11,460 acres of openings (non-fisher habitat) were calculated and these included openings of Forest Service land, but the majority were present on private property.

Upon analysis of the existing condition, this analysis area is below the desired thresholds, due to a large presence of non-Forest Service lands and past activities. Total mature habitat represents about 13%, well below the 50% mature habitat (Sauder 2014, Sauder and Rachlow 2014). Open habitat contributes to a 39% fragmentation level which is above the 5% open areas (Sauder 2014, Sauder and Rachlow 2014). In summary, the potential territory analyzed for the fisher is unlikely to meet the thresholds suggested by researchers because of the large presence of private land that is not held to the same guidelines as those for the forest.

Population Trends: Fishers have a global ranking of G5 (global rank for demonstrably widespread, abundant and secure) and a state rank of S2 (imperiled), (Nature Serve 2019).

## **Gray Wolf**

Gray wolf populations were extirpated from the western U.S. around the 1930s. Over time, individual wolves from Canada occasionally dispersed into Idaho. The gray wolf was listed as an endangered species in 1978. In the mid-1990s, gray wolves were introduced into central Idaho. By 2011 the USFWS finalized the delisting of the wolf in Idaho (IDFG and Nez Perce Tribe 2014).

Wolf habitat spans a broad range of elevations and habitat types. Key habitat components include: 1) a sufficient year-round prey base of ungulates and alternate prey; 2) suitable somewhat secluded denning and rendezvous sites; and 3) sufficient space with minimal exposure to humans (USDI FWS 1987). Denning/rendezvous sites, elk habitat effectiveness, and elk security areas (see Elk section) are used to assess existing conditions for wolves. Maintaining elk habitat effectiveness above minimum Forest Plan standards, providing elk security areas above minimum recommendations, and managing winter range to enhance forage productivity and quality would provide a sufficient prey base to sustain wolf populations at State objectives.

Population Trends: The gray wolf has a global ranking of G5 (global rank for demonstrably widespread, abundant and secure) and a state rank of S4 (apparently secure), (Nature Serve 2019). The wolf population on the Forest is considered as stable to increasing.

## **Long-eared and Long-legged Myotis**

Both bat species typically roost in snags, rock crevices, and caves. In forested areas, they have been found in snags and exfoliating tree bark. The long-legged myotis is more closely associated with coniferous forest habitat than the long-eared myotis. Both bats are known to forage together (Johnson, Lacki, & Baker, 2007).

The long-eared bat has been found roosting in the snags and stumps of Douglas-fir, western hemlock (Barclay & Kurta, 2007), western red cedar (Arnett & Hayes, 2009), and pine (Vonhof and Barclay, 1997)

Long-legged myotis are medium-sized bats that prefer large snags for roosting, but will also roost in live trees. Arnett and Hayes (2009) found that long-legged myotis infrequently roosted in snags or trees in stands less than 40 years old, and 58% of the snag roosts and 33% of the live tree roosts were located within riparian management buffers retained during harvest near small- and medium-sized perennial streams. Lacki et al (2010) also found that retention of habitat in riparian areas was important. Long-legged myotis roosted in snags in mid-seral (41-80 years) and old growth stands.

Population Trends: Nature Serve (2019) ranks the long-eared bat as G5- secure globally, and the long-legged bat as G4- apparently secure. Both species are considered in Idaho at the state rank of S3- rare or uncommon, but not imperiled.

## **Western Toad**

Toads breed in temporary and permanent lakes, ponds, streams, and road ditches. They prefer shallow, warm areas with mud bottoms, and typically breed in May and June. Potential breeding and dispersal habitat occurs throughout the area along the network of riparian areas. Toads can be found from dry grasslands to moist subalpine forests, but optimal habitat is found in humid areas with moderate undergrowth. They are largely terrestrial, but generally found within fair proximity to water.

Adult western toads are largely terrestrial and are very active at night. They have been known to move up to 1 mile from their breeding habitats, (Bartelt et al. 2004) often into upland habitats (Bull 2006). Toads selected south-facing slopes, preferred open sites to forested settings, and sites with high density of burrows, rocks, logs, or rootwads that provided cover (Bull 2006). Burned and harvested sites were not avoided by western toads in Bull's study. Guscio et al. (2007) found western toad occurrence increased after wildfires and they used severely burned areas. Use shifted from severely burned to moderately burned areas in the late summer likely as a result of more ground/canopy cover and higher soil moistures.

Population Trends: The toad has a regional rank of G4 (apparently secure), and for Idaho the state rank is S2: imperiled (Nature Serve, 2018). Declines in abundance have been reported throughout the species' range due to disease and parasites.

## Neotropical Migratory Birds

Neotropical migratory birds are species that breed and rear their young in the United States and Canada, then migrate south to winter in Mexico, the Caribbean Islands, and Central and South America. Under the National Forest Management Act (NFMA), the Forest Service is directed to “provide for diversity of plant and animal communities based on the suitability and capability of the specific land area in order to meet overall multiple-use objectives” (P.L. 94-588, Sec 6 (g) (3) (B)). The January followed by the US Shorebird Conservation Plan and Executive Order 13186 (USA, 2001), and the January 2004 PIF North American Landbird Conservation Plan (Rich et al., 2007) all reference goals and objectives for integrating bird conservation into forest management and planning.

In late 2008, a Memorandum of Understanding between the USDA Forest Service and the US Fish and Wildlife Service to Promote the Conservation of Migratory Birds was signed. The intent of the MOU is to strengthen migratory bird conservation through enhanced collaboration and cooperation between the Forest Service and the Fish and Wildlife Service as well as other federal, state, tribal and local governments. Within the National Forests, conservation of migratory birds focuses on providing a diversity of habitat conditions at multiple spatial scales and ensuring that bird conservation is addressed when planning for land management activities.

In the absence of activities or natural disturbances, the forest would continue grow towards a more closed canopy. Avian species that benefit are those that thrive in such habitats include those that nest and forage in the tree canopies: insect gleaners such as woodpeckers, some warblers, vireos, and seed eaters such as grosbeaks. For some neotropical migrants the project area may not provide the desired habitat, which usually results in the species moving on.

## American Marten

Marten is added to the analysis as it was inadvertently missed while preparing the preliminary environmental assessment.

In the Idaho Panhandle and the Nez Perce-Clearwater Forests, tree species in mature mesic forests associated with marten habitat include western red cedar, subalpine fir, Engelmann spruce, western larch, and lodgepole pine (Wasserman et al. 2010; Koehler and Hornocker, 1977; Koehler et al. 1975). Marten have also been seen in ponderosa pine stands. Some studies have found that marten avoid open areas such as clearcuts or other fragmented habitats (Shirk et al. 2014; Wasserman et al. 2012; Moriarty et al. 2011; Potvin et al. 2000; Hargis et al. 1999; Buskirk and Ruggiero 1994; Koehler et al. 1975).

Many research articles (Shirk et al. 2014, Wasserman et al. 2010, Hargis et al. 1999, Buskirk and Ruggiero 1994, Koehler 1975) identified potential marten habitat associated with cool, moist habitat types with the following tree species: subalpine fir, spruce, lodgepole pine, grand fir, and western red cedar. Another attribute used for marten habitat was tree canopy cover had to be 30% or greater based on some of the previous research, as well as Bull et al. 2005, and Snyder and Bissonette 1987. Marten habitat is well connected by large contiguous blocks with high canopy closure, riparian areas and old growth. The median size of the largest forest patch in marten home ranges was 150 ha (370 ac) for females and 247 ha (610 ac) for males (Chapin et al. 1998); 3,187 ac for male, 2,495 ac for female (Shirk et al. 2014).

The project area appears large enough in size to support a marten territory, however the potential habitat is lacking. Potential habitat in the project area is small (45 acres) and distributed in 3

disjointed patches. The project area is under-represented or lacking sub-alpine fir, Engelmann spruce, and lodgepole; and has about 30 acres of western red cedar. Dominant tree species representation is grand fir and shade tolerant mix of about 735 acres that may be used by marten, but the project area is generally lacking the tree species composition as mentioned by the above researchers.

Population Trends: The marten has a regional rank of G5 (secure), and for Idaho the state rank is S5: secure (Nature Serve, 2019). Declines in abundance have been reported throughout the species' range due to disease and parasites.

## **Northern Goshawk**

The northern goshawk was identified as a Forest MIS for old-growth forest. Goshawks use large landscapes, integrating a diversity of vegetation types over several spatial scales to meet their life-cycle needs (John R. Squires & Kennedy, 2006). In "The Northern Goshawk Status Review," the USFWS found that the goshawk typically uses mature forest or larger trees for nesting habitat; however, it is considered a forest habitat generalist at larger spatial scales (USDIFWS, 1998). The FWS found no evidence that the goshawk is dependent on large, unbroken tracts of "old growth or mature forest."

Goshawks prey in habitats that contain snags, downed logs, woody debris, large trees, herbaceous and shrubby understories, and a mixture of stand structural stages (Wisdom et al., 2000).

Foraging habitat for goshawks may occur along the edges of open areas; such as meadows, burned areas, timber units, streams and roads. Forage habitat is not considered limiting factor for the raptor on this Forest.

A recent study estimated male breeding-season home range at 5,146 acres and the female range at 3,859 acres. (Moser and Garton 2009). A 5,000-acre area (about the size of a male breeding home range) was calculated that includes the project area. Potential goshawk nesting habitat was queried at 1,808 acres within the potential breeding range.

Nest areas are usually mature forest with large trees, relatively closed canopies (60-90%) and open understories (Squires and Kennedy, 2006). In central Idaho, goshawks nest in a variety of forest stands that are comprised of mature trees with relatively high canopy cover and open understories (Moser, 2007). Favored habitats typically are located in forest stands having only 1 or 2 canopy levels with an open or mixed-density understory (ibid). Another study found that nesting areas have multiple alternative nests (Moser and Garton 2009). The researchers found the average size of a nesting area in their Idaho studies were around 420 acres (Moser and Garton 2009, Moser 2007).

Nesting habitat was chosen indicator as it is the primary limiting factor for goshawks and is represented by a much narrower range of vegetation structure and composition than the post-fledgling areas and forage area.

Home ranges are likely not defended from other goshawks, with the exceptions of the nest area and post-fledging area (Brewer et al. 2009). Home ranges of adjacent pairs may overlap (Squires & Reynolds 1997; Squires and Kennedy, 2006). Goshawks have been found to use the same nesting area for decades, and goshawk territories typically contain a number of alternate nests (Moser 2007). Goshawks appear to range over large areas and use a variety of habitats outside of the nesting area (Kennedy 2003).

Population Trends: The goshawk is rated secure across its range (global rank G5) and S3 in Idaho; which is rare or uncommon, but not imperiled (Nature Serve 2019).

## **Pileated Woodpecker**

The pileated woodpecker is another Clearwater National Forest MIS for old growth forest and large snag habitat. Pileated woodpeckers are large, cavity-nesting birds associated with late successional stage forests, but also may use younger forests that have scattered, large, dead trees (Bull & Jackson, 1995). The woodpecker is common in both cut and uncut mid-elevation forests and appear to do well in a matrix of forest types (Hutto, 1995). One group of researchers conducted a study on the density of pileated woodpecker nesting pairs in areas before and after timber harvest activities (Bull et al. 2007). In six of the seven study areas, the density of the nesting pairs were unchanged, or increased or decreased by only one pair. In these sampled areas, the amount of mature and old forests decreased by less than 25%, with consistent pileated woodpecker densities.

Feeding habitat for pileated woodpeckers is highly dependent on the availability of carpenter ants which make up the majority of their food supply (Aney & McClelland 1990). Preferred feeding habitats have high densities of snags and logs, dense canopies, and tall ground cover, with more than 10% of the ground area covered by logs.

The mean size of nest trees ranged from 28" dbh in Montana (Aney and McClelland 1990) to 33 inches dbh in Oregon (Bull and Jackson, 1995). The minimum canopy cover selected by pileated woodpeckers for nesting stands ranges from 15 to 60 percent depending on the habitat type (Bull et al. 1992), (Bull & Holthausen 1993); (Bonar, 2001).

Territories of nesting pairs cover 500-1000 acres in Montana, 1000-1300 acres in western Oregon, 320-600 acres in northeastern Oregon (Aney and McClelland 1990), and about 1,000 acres in another Oregon study (Bull and Holthausen, 1993). The nesting territory of 1,000 acres (Bull et al. 2007) has been used for this analysis because this study on the woodpecker has been conducted for at least 30 years, in habitats similar to those on this forest.

The current condition of nesting habitat in the territory (86 acres) is considered the most limiting factor for pileated woodpeckers. Nesting habitat is a more specialized range of vegetation structure and composition than the stand age and structure for foraging habitat. The nest tree is the most important variable to estimate breeding habitat use by the pileated woodpecker (Kirk & Naylor 1996; Giese & Cuthbert 2003).

Not every stand within a bird's home range is used as feeding habitat, and the range of a nesting pair is partly determined by the amount of suitable feeding habitat in proximity to the nest site.

Population Trends: The pileated woodpecker is rated secure across its range (global rank G5) and apparently secure (state rank S4) in the state of Idaho (Nature Serve, 2019).

## **Rocky Mountain Elk**

Elk is a management indicator species for commonly hunted big game species, and an MIS for general forest seral species easily affected by management activities on the Clearwater National Forest.

Elk are habitat generalists and use a diversity of forest types and structures that provide forage and hiding cover (Unsworth et al. 1998). They forage in meadows and early seral communities

from spring through early summer, use more closed canopies from late summer through fall, and rely upon low elevation, warm aspect, and snow-free or snow-limited areas for foraging in the winter. Adult bulls often winter at much higher elevations than cows and immature elk. Elk also require forest cover for security and thermal regulation (Ward et al. 1979), although the results of (Cook et al. 1998) did not find that such forest cover significantly enhanced the condition of elk in winter or summer, and it did not assess effects of topographic or other landscape features that could enhance energy conservation by protecting from wind or enhancing absorption of solar radiation.

The information presented during the combined scoping/30-day comment period discussed recent literature on elk that is not evaluated or analyzed by the Servheen et al. 1997 guidelines required by the Forest Plan. Discussion of literature that does not pertain to the elements of elk analysis would not be evaluated.

Roads built into elk habitat increase hunter access, increasing elk vulnerability to harvest (Unsworth et al. 1993; Christensen et al. 1993). Other literature on elk modelling has suggested buffers for road effects (Ranglack et al. 2017; Frair et al, 2008; Rowland et al. 2005; Rowland et al. 2000), size of elk unit to be analyzed (Rowland et al, 2005; Boyce et al, 2003; Unsworth et al. 1998; Christensen et al. 1993), the influence of different road types on elk space use across seasons and by sex (Montgomery et al. 2012), and other elk habitat considerations. Field observations by the biologist, other forest personnel and public individuals show that elk may use roads as evidenced by elk tracks seen on roads. In some cases creation of temporary roads on ridges may be used by elk as game trails.

### **Stray Creek Existing condition**

The previous description of elk habitat discussed recent literature concerning factors that influence elk recruitment. Project effects on elk will follow the current Forest Plan direction that is to use the elk analysis developed by Servheen et al. 1997. The forest plan standard for the affected elk analysis area (EAA) is for elk habitat effectiveness to equal or greater than 25% elk habitat effectiveness (EHE). This effectiveness is calculated by a combination of variables that may affect elk in the area and that is the amount of openings within forest, security areas, hiding cover, motorized vehicle use periods, and livestock presence. All these variables combine to create a value for EHE. Currently, the one EAA that encompasses the project area is meeting forest plan standards (EHE is 47%).

### **Shiras Moose**

The moose is a forest MIS species for mature timber with understories of Pacific yew. The latter habitat is considered primary winter habitat for the ungulate (Pierce and Peek 1984). Other preferred forage (besides yew) of moose include false huckleberry, Scouler willow, serviceberry and alder. Research in the Canadian Rockies found moose browsed lodgepole pine and subalpine fir (Poole and Stuart-Smith 2005). During late spring to fall moose have been seen in shrubfields, riparian areas and locations recently burned or timber harvest units.

Pacific yew was declining in the early 1990s due to the discovery of taxol, an anticancer agent found in the tissues of the plant (Busing and Spies 1995). In anticipation of the increased demand and harvest of the plant, the Forest Service implemented interim management guidelines for the conservation of yew in 1992. The yew is a slow-growing plant that is most abundant mature to old-growth forests.

Moose is a big-game species found in relatively low numbers, scattered across the Forest with the exception of the Powell Ranger District. Near this Idaho/Montana boundary, moose occur in larger concentrations (USDA Forest Service 1987). A statewide report documents that moose have expanded in their range and numbers in parts of Idaho (IDFG 2017). However, the same report notes that moose have been declining in portions of the Clearwater region in the past 15-20 years. Some reasons the study offered were moose tags have been reduced, and moose range has increased in areas once thought to be less than optimal habitat.

The project area is located in the eastern portion of GMU 10A. The management direction for the game management unit is to allow moose populations to increase where habitat and other conditions will support expansion, moose harvest will be adjusted according to hunter success rates and known mortalities will be documented (IDFG 2016). The IDFG manages hunting pressure by controlled hunts, which disperses hunters and/or directs harvest in specific areas. The agency finds the expansion of wolves on moose populations are as yet undetermined, and other factors- disease, parasites and nutritional data are still being evaluated across the state.

Population Trends: The moose is rated as G5 (abundant and secure in the U.S. and Canada), and S5 in Idaho: abundant and secure (Nature Serve 2017). A recent publication estimates around 10,000 moose in Idaho (Nadeau et al. 2017).

## Environmental Impacts

### No Action

In general, the no action alternative would not create disturbance or loss of habitat by human actions. Wildlife species that prefer open areas may experience the loss or decrease of habitat from tree succession. Species that prefer more closed forest or canopy cover are likely to benefit from tree succession. If no action, natural processes would continue such as wildfires, windstorms, insect and disease outbreaks, and other weather events may impact the habitats of the wildlife species analyzed below. Habitat would persist as described in the existing condition detailed above.

### Proposed Action

#### Sensitive Species

##### *Fisher*

The fisher is an Idaho species of greatest conservation need (IDFG 2018). One record of a fisher trapped and released in the Stray Creek project area occurred in 1989.

Sauder (2014) conducted fisher research on the Nez Perce-Clearwater National Forests and estimated the average home range for a male fisher is about 24,300 acres and 12,200 acres for a female. Such acreage is achieved by analysis at the HUC 12 (hydrologic unit compartment) scale which is the area of analysis for the predator. The Middle Lolo Creek HUC encompasses about 29,520 acres, including the project area. The analysis area straddles about 10,020 acres of Forest Service ownership, while the other 19,500 acres are under private ownership.

The model used for analyzing fisher habitat was developed by Sauder but does not include private lands. Potential habitat (patches of dense forest) was interpreted from a NAIP satellite photo of the on private lands. The combination of the model and satellite imagery presented about 18,053

acres (61%) of potential fisher habitat in the analysis area. The amount of potential habitat suggests that the area would support one female territory.

Further analysis of open areas resulted in 11,467 acres (39%) was considered as non-habitat and lacking forest or dense forest cover for the fisher. Total mature fisher habitat was calculated at 13% representation in the HUC. This is below the recommended 50% representation in a territory (Sauder 2014; Sauder and Rachlow 2014). General fisher habitat (modelled) represents 19%. Open or non-fisher habitat is at 39%, well above the 5% threshold recommended by the authors noted above. Most of the private lands (57%) have undergone timber harvest or other disturbances that are not supporting potential fisher habitat. Additionally, the privately-owned portion of the analysis area is highly fragmented with roads, clearings and recent harvests. As mentioned in the Affected Environment section, this potential territory is not supporting the desired conditions as mentioned by the above researchers. Additionally, it is not likely to provide the desired levels of mature habitat or low fragmentation due to the amount of private property.

### **Direct and Indirect Effects**

The proposed project would harvest 297 acres of general habitat and 128 acres of mature fisher habitat. At the analysis scale, the activities would reduce mature habitat to 12.8% and general habitat to 18.2%, including 8 acres affected by temporary road construction. Openings would increase to 40%. No harvest would occur in old growth.

During project implementation, a fisher may be disturbed or displaced by the harvest activities (noise and movement by man and machine). The timber harvest would create openings that the fisher would likely avoid. However, the predator may forage along the edge of the units and undisturbed forest.

### **Cumulative Effects**

The cumulative effects boundary is the analysis area described above. The cumulative effects timeframe is over 40 years; as the period for regeneration harvest areas to develop into mature habitat of 10 inches dbh or greater (as defined for fisher habitat by Sauder and Rachlow, 2014).

Current/on-going activities authorized in the in Lolo Insect & Disease project would affect about 85 acres of fisher habitat. The associated harvest would begin the year prior to implementation of the Stray Creek activities. The combined reduction from the proposed action and the Lolo Insect & Disease project would reduce fisher general habitat to 18%, and mature habitat to 12.6%. This would maintain less than the 50% representation suggested by Sauder 2014 and Sauder and Rachlow 2014. Habitat fragmentation from the combined projects would be about 40.5% which is above the 5% threshold recommended by researchers (Sauder 2014, Sauder and Rachlow 2015). Disturbance and displacement of an individual fisher is also possible. Additionally, slash burning, tree planting, road building and decommissioning, fire suppression, and wood cutting are other activities that may perturb a fisher near such activities.

No harvest would occur in old growth or riparian areas, which would retain resting and denning habitat, as well as movement corridors for the fisher. The cumulative effects of the proposed activities would decrease habitat and raise fragmentation levels in this potential territory, which may be undesirable to an individual fisher. Additionally, the majority of the analysis area is in private ownership, which is not bound to regard the best interest of fishers. If all the Forest Service land provided fisher habitat, fragmentation would still be at 37%, which is above the thresholds (Sauder 2014, Sauder and Rachlow 2015).

## **Determination**

Areas affected by project activities (harvest and roads) would provide at least 10% cover in a fisher territory; about 10-15 years post project completion. Forty years after the project is completed, mature forest would return to the affected areas, unless other disturbances (man-made or natural events) occur.

The noise and movement by man and machine, as well as removal of fisher habitat would likely displace an individual fisher from the project area. Human presence would continue for up to 5 years of additional entries in the harvested unit for fuels treatments, tree planting and road/trail obliteration. Contiguous fisher habitat is adjacent to the project area and provides displacement habitat for an individual to retreat to. A recent Forest-wide query calculated over one million acres of fisher habitat on the Clearwater National Forest; about 505,000 acres are mature habitat. Thereby, the forest would still maintain habitat for territories, forage, and shelter for the mammal. The effects of the Stray Creek Project may impact individuals or habitat but will not likely contribute to a trend towards federal listing or cause a loss of viability to the population or species of fisher.

## ***Gray Wolf***

After the wolf was de-listed in Idaho, 2011, population control plans on the predator has intensified. Two major issues surrounding wolf management are depredations on livestock, and impact of predation on elk, moose, and other ungulates (IDFG 2017). The goal of the state management agency is to meet the intent of the 2002 IDFG Wolf Plan: maintain about 15 wolf packs in the state. Wolf zones are no longer used as an analysis tool. Instead, incidents of livestock depredations are tracked by Game Management Units or GMUs. The project area is located in a portion of GMU 10a (Dworshak Elk Zone). Currently, this unit is considered at a moderate intensity of predation on Idaho elk populations identified in the IDFG elk management plan of 2014 (IDFG 2017). The wolf population on the Forest is considered as stable to increasing.

## **Direct and Indirect Effects**

The proposed action would not affect wolf habitat. Implementation of timber harvest operations (noise and movement by man and machine) may temporarily agitate or displace a wolf from the project area during periods of activity. Wolves are known to return to harvested units during hours of darkness to hunt for prey. The project area is part of a larger elk analysis area, which meets the Forest Plan guidelines for elk habitat effectiveness (see elk section).

## **Cumulative Effects**

The effects area is the Dworshak GMU (318 square miles). The GMU is large enough to provide year-round prey, secluded denning and rendezvous sites, and secluded areas from human contact. The time frame for a potential increase of big game in this area is about 15 years post after harvest operations.

The Stray Creek Project would create about 425 acres of openings in tree habitat that would provide forage for big game (elk, deer, moose); which are important prey for the wolf. The early forest vegetation would provide forage for elk and other ungulates from 2-15 years. As the big game return, the wolf would likely visit these areas for potential prey. Seedlings will be planted and would offer hiding cover for elk in a span of 10-20 years. This cover in affected units would reduce detection of elk along open roads during hunting season.

A portion of the Lolo Insect and Disease project is in the same GMU as Stray Creek, and proposes to harvest about 4.4% of the land in the GMU analyzed. This will provide more forage for big game, and increase the prey base for the wolf. During implementation wolves may be temporarily disturbed or displaced, but they will return when the prey base is available.

Firewood cutting is likely to occur along roads open to public motorized access, and may temporarily displace a wolf near the area of activity. Fire suppression would continue which would displace wolves in the area of operations. Again, forest openings created by fire would produce future forage for big game and offer the wolf more opportunities for prey.

### **Determination**

Control of wolves would continue until the state has reached the objective of 15 wolf packs. The effects of the Stray Creek Project may impact individuals or habitat, but will not likely contribute to a trend towards federal listing or cause a loss of viability to the population or species of the gray wolf. The wolf status is considered as stable to increasing on the Forest (NP\_CNF Monitoring Report 2018).

### ***Long-eared and Long-legged Myotis***

Modelled habitat in the project area is estimated at 374 acres (trees of the size of 15 inches or greater).

### **Direct and Indirect Effects**

Timber harvest is proposed in 134 acres (less than 36%) of potential bat habitat. None of this would occur in old growth or riparian areas. The retention of riparian areas is an important habitat feature as well (Lacki et al. 2010). The regeneration harvests would retain a few large trees or snags that may provide roost habitat for the bats. However, the harvest activity would reduce habitat in the project area. The action alternative would disturb, displace or maim individual bats in the proposed units. If harvest occurs during the winter season, bats would not be affected as they would be present in their wintering roosts in the southern part of the state.

Implementation of the proposed action would occur over a period of approximately 10 years; thereby, not all units would be treated at once, and not all activities would occur at the same time. Treatments would reduce tree canopy cover, and reduce potential habitat. As the understory recovers, forbs and shrubs would increase and offer habitat for insects and bugs. Plants that produce flowers would attract butterflies and moths, while the dead or dying trees would provide habitat for beetles and other bugs. Both butterflies and beetles are the preferred staple of the long-eared and long-legged bat's diet (Lacki et al. 2007). The pulse in bat forage would last until the tree canopy shades out the understory; about ten to fifteen years.

### **Cumulative Effects**

The project area is the analysis area, as it is large enough to host many bats for the summer migration season. The time period relevant to bat presence in the analysis area is late spring to early autumn: the period of their migration for richer food sources.

Firewood cutting would remain limited to the low mileage of road open to the public; which remains the same as the existing condition. Fire suppression would continue, which may reduce the loss of bat habitat to wildfire.

## **Determination**

About 240 acres of potential bat habitat would remain in the project area. Another 150 acres of potential habitat is adjacent to the project area; which is available for displaced bats. The proposed action may impact individuals or habitat, but will not likely contribute to a trend towards federal listing or cause a loss of viability to the population or species of the analyzed myotis species. Harvest prescriptions would retain large trees and snags within units, and design criteria avoids effects to habitat by coordinating prescribed burn efforts to reduce disturbance and retain trees with cavities and/or large nests.

## ***Western Toad***

Breeding and larval stage development occurs in water: ponds, puddles, slow-moving streams and so on. About 181 acres of such breeding habitat is available in the project area during spring and early summer. Upland habitat in the rest of the project area comprises 658 acres. Timeframe for the effects analysis (direct, indirect, and cumulative) is approximately 10 years.

## **Direct and Indirect Effects**

No project activities would occur in riparian areas. Therefore, breeding and development of young larvae to adults would occur without loss or disturbance of habitat in the moister habitats. About 425 acres of upland habitat would be affected by timber harvest. All activities have the potential to disturb, displace, maim or kill an individual toad outside of a riparian zone during project implementation.

About 51% of upland habitat for the toad in the project area would be affected by timber harvest. Downed woody material would increase exponentially, which provides hiding and burrowing cover for adult toads, as well as other invertebrates that the amphibian preys on.

## **Cumulative Effects**

Fire suppression would continue to be implemented if potential harmful effects are anticipated to human-owned property. There is potential for no cumulative effects because there are no future foreseeable activities to analyze.

## **Determination**

The proposed action may impact individuals or habitat, but will not likely contribute to a trend towards federal listing or cause a loss of viability to the population or species of the western toad. Harvest prescriptions would retain downed wood and snags (future downed wood) within units, and design criteria avoids effects to potential breeding habitat (all riparian areas).

## ***Neotropical Migratory Birds***

The project area is 98 percent forested: therefore, migratory birds that prefer open areas without trees would have about 10 acres of available habitat. More open habitats are available on private or state-owned holdings, to the west of the project area. Ponderosa pine and Douglas-fir occupy about 78 acres in the project area. This habitat may support migrants such as the flammulated owl, American robin, and western tanager. Closed forest of grand fir and western red cedar dominate about 750 acres in the project area. The denser forest would support thrushes, kinglets, and some warblers.

## **Direct and Indirect Effects**

In the absence of activities or natural disturbances, the forest would continue grow towards a more closed canopy. Avian species that benefit are those that thrive in such habitats include those that nest and forage in the tree canopies: insect gleaners such as woodpeckers, some warblers, vireos, and seed eaters such as grosbeaks. For some neotropical migrants the project area may not provide the desired habitat, which usually results in the species moving on.

Proposed harvest activities would occur in 425 acres. Temporary roads would reduce less than one acre of forest habitat. No harvest activities or temporary roads would occur in riparian areas. Newly harvested or burnt areas would provide habitat for migrant species that prefer open habitats. Conversely, this change would not be favorable to avian species that prefer closed habitats. The latter would move to adjacent areas of forest within or adjacent to the affected areas. Harvested units would recover from native seed sources in the soil and planted trees. During the first 15 years after timber sale completions, growing shrubs and trees would offer good opportunities for nesting songbirds. A greater quantity and diversity of invertebrates would be available in this life stage, which would benefit bird insectivores. As tree densities increase, the bird species that prefer closed habitats would find these areas more suitable for nesting habitat. Noise and movement of machinery and other human activity may disturb migrant birds. The operating season may disrupt some nesting birds in or near areas of project activities. Harvest operations would begin in July, when some bird species may have fledglings present near their nest. However, most of the timber harvest and prescribed burning would occur after young birds have left the nest. No activities would occur in old growth.

Timber-harvested units would be restocked by tree planting, while burned areas would recover from native seed sources. During the first 15 years after timber sale completions, growing shrubs and trees would offer good opportunities for nesting songbirds. A greater quantity and diversity of invertebrates would be available in this life stage, which would benefit bird insectivores. Proposed activities that reduce vegetation would be beneficial to bird species that prefer more open habitats (shrubfields, small meadows or open fields). Species that would benefit include hummingbirds, flycatchers, red-tailed hawk, sparrows, and some ground-nesting birds.

## **Cumulative Effects**

The cumulative effects area is the project area because it is large enough to offer habitat for a breeding cycle of forest-preferring migratory bird species. The proposed actions (temporary road construction and obliteration, timber harvest, burning, tree planting) would be completed in approximately 10 years. Fire suppression may occur for wildfires that may break out in the area. A wildfire and suppression efforts may displace or harm an individual migratory bird. However, most fires occur after the nesting season, and the new generation of young birds would have the ability to fly away from danger posed by a fire or suppression efforts.

Long-term effects would be the benefit of increased vegetation for forest preferring migratory birds. Maintenance of vegetation to open ponderosa pine stands would benefit the migratory flammulated owl. No new permanent openings, such as roads would occur in the project area. Temporary openings created by harvest may encourage predators (ravens, starlings) to compete with neotropical migrants. Additionally, cowbirds may be attracted by the edge-effect habitat of units to seek "host nests" for parasitism of neotropical migrants. Temporary roads would be decommissioned after use, and would provide potential habitat (vegetation) for migratory birds. New vegetation would offer nectar (flowers) or invertebrates that live or feed on the plants.

Flower nectar would provide food for hummingbirds, while an increase in insects. There is potential for no cumulative effects because there are no future foreseeable activities to analyze.

### **Determination**

The proposed action may impact individuals or their habitat, but is not expected to result in a loss of viability in the planning area, nor cause a trend toward federal listing of migratory birds, because the short period of timber operations, and design criteria (WL-1, and 2) would maintain potential habitat and minimize disturbance during the period of migratory bird presence in the area.

### **Management Indicator Species**

#### ***American Marten***

The American marten is a Clearwater Forest MIS for mid-high elevation, mature forests. Habitat in the project area is small, due to the lower elevations present in the area, and many of the tree species associated with marten habitat is under-represented in the project area.

### **Direct and Indirect Effects**

The analysis area is the project area, and the time frame encompasses the 10-year period of potential activities. No old growth would be harvested in the project area. The proposed action would harvest 35 acres. No temporary roads would occur in marten habitat.

### **Cumulative Effects**

The analysis area is the project area, and the time frame encompasses the 10-year period of potential activities. Past harvest (1965-2011) affected about 516 acres: only 25 acres were impacted by regeneration timber harvest (see Forest Vegetation section of the EA). About 200 acres of potential marten habitat would be available adjacent to the project area for displacement. Wildfires may impact marten habitat in the project area. Fire suppression may retain some of this habitat, however, the amount of habitat suggested for a marten territory would not occur from such action. There is potential for no cumulative effects because there are no future foreseeable activities to analyze.

### **Conclusion**

The proposed project activities may disturb or displace an individual marten. The area is lacking potential habitat (cool, moist habitat types) for marten because the tree species composition (sub-alpine fir, Engelmann spruce, and lodgepole) does not occur. Additionally, over 1 million acres of potential habitat was documented for the marten, and its status is considered stable (NP\_CNF Monitoring Report 2018).

#### ***Northern Goshawk***

The northern goshawk is a management indicator species for old growth. Regional guidelines suggest that goshawk analysis occurs at the 5,000-acre breeding range (Brewer et al. 2009). About 1,808 acres (including old growth) provides nesting habitat for the raptor. The arrangement of the habitat is mostly contiguous. Moser and Garton (2009) suggested that potential nesting areas to be around 420 acres. The timeframe for effects would be 80 or more years for nesting habitat in the timber harvest affected areas.

### **Direct and Indirect Effects**

No old growth would be harvested in the project area. The proposed action would harvest about 121 acres of potential nesting habitat. The construction of new temporary roads would affect less than one acre of habitat. The proposed action would reduce potential nesting habitat by 7% in the analysis area. Nesting habitat would remain present in or adjacent to all nesting territories. Noise from man and machine, and the increased openings created for harvest would likely displace a goshawk in the area. If an occupied nest is detected, then activities would be suspended during the nesting period. Large trees and other forest habitat that is retained, would continue to provide foraging areas for the raptor. Over 1,600 acres of contiguous nesting habitat would be available within and along the western boundary of the analysis area as potential displacement nesting area.

Project activities may disturb a nesting pair of goshawks. Design criteria would protect a nest (WL-2 and 3) and prohibit activities in an active post-fledgling area (WL-4). Researchers found that there were no treatment effects on "goshawk re-occupancy, nesting success, or number of fledglings between harvested and unharvested nesting areas" (Moser and Garton 2009). Additionally, the authors found that goshawks have alternative nests in a territory that may be used if a nest is lost.

### **Cumulative Effects**

Fire suppression would occur in the event of a wildfire within the project area that may displace a goshawk. There is potential for no cumulative effects because there are no future foreseeable activities to analyze, this includes no effects expected from the Lolo Insect & Disease project because no units occur (spatially) within the analysis area.

### **Conclusion**

The proposed project activities may disturb or displace an individual goshawk. No old growth would be removed. Potential nesting habitat is available and adjacent to the project area. Over 52,000 acres of nesting habitat was documented for the goshawk and its status is considered stable (NP\_CNF Monitoring Report 2018).

### ***Pileated Woodpecker***

The pileated woodpecker is a management indicator species for old growth and large snag habitat. Bull et al. 2007 estimate the woodpecker uses a home range of about 1,000 acres in size. The potential nesting habitat within this home range was calculated at 127 acres. The effects timeframe is 80+ years, as it would take this long for regeneration harvest areas to develop into mature stage and later- old growth habitat in about 150 years.

### **Direct and Indirect Effects**

The proposed action would reduce about 5 acres of nesting habitat for the woodpecker. The affected acreage for the woodpecker is spread out across two patches of habitat. This would reduce the woodpecker's habitat in the proposed action by 7% in the in the home range that was analyzed. Some large tree habitat, as well as snags would be retained for woodpecker foraging habitat. Project activities may disturb a nesting pair of woodpeckers. About 68 acres of nesting habitat is near the project area which would provide habitat for a woodpecker and would remain in the home range; which provides habitat for a woodpecker displaced by project activities. Design criteria (WL-2) would protect a discovered nest.

## **Cumulative Effects**

Fire suppression would occur in the event of a wildfire within the project area that may displace a pileated woodpecker. There is potential for no cumulative effects because there are no future foreseeable activities to analyze, this includes no effects expected from the Lolo Insect & Disease project because no units occur (spatially) within the analysis area.

## **Conclusion**

The proposed project activities may impact habitat and disturb or displace an individual. Snags will be retained in treated areas according to the Forest Plan by implementing the silvicultural prescription. Displacement habitat is available and contiguous with habitat that may be affected by the proposed activities. Over 142,000 acres of nesting habitat was documented for the woodpecker, and its status is considered stable (NP CNF Monitoring Report 2018).

## ***Rocky Mountain Elk***

The elk is a management indicatory species for big game habitats. The project area is located in summer elk range, and in a portion of one elk analysis area (EAA). Forest Plan guidelines (Servheen et al. 1997) for the affected Yakus EAA is to maintain at least 25% elk habitat effectiveness (EHE) (Forest-wide standard 7.j.(3)). The EHE calculation is a sum of calculations based on the following: size of the EAA; size of certain areas for security, openings, hiding cover, road densities, elk use related to cattle density, and size and distribution of cover and forage. As a project is implemented, EHE may be affected by changes in the above metrics. Calculations for each EAA during each phase of the activities are available in the project record.

## **Direct and Indirect Effects**

The elk analysis for the Yakus EAA was updated that utilized a more accurate road layer than the previous analysis. More roads were displayed in the area by the more recent road information. This finding yielded an increase in road densities, which decreased the EHE in the elk analysis area.

The existing condition of EHE is 40% for the Yakus EAA. The proposed action would increase openings in the EAA due to timber harvest,

Such actions reduce hiding cover and security areas Temporary roads and harvest treatments along system roads create or increase forest openings. Vegetation recovers naturally or from re-stocking the open areas with trees. As vegetation recovers; elk hiding cover and security would return.

Elk habitat effectiveness (EHE) would decrease to 38% during the proposed action activities. Prescribed burns would reduce fuels in the understory, and minimally affect hiding cover in the short-term. This treatment would encourage growth of a new understory of plants and shrubs for big game forage. Vegetation in the harvest units and along the open road would provide hiding cover for elk in about 10-15 years after the project is completed.

After the proposed action is completed, road densities usually drop, due to closing and/or decommissioning roads used in the harvest operations. Road closures usually increase elk security. Openings along roads also affect road densities as a road section that is along an opening is evaluated at a higher co-efficient than a road bordered by vegetation. The post action EHE value does not reach that of the existing condition due to the increased openings along a road that is open all year. This factors in as a higher road density post action because these roads provide a

higher vulnerability for elk during hunting seasons and that is why the Yakus EAA has a lower EHE value after the project is completed, as opposed to the existing condition. Table 2 tracks the changes in the EAA located in this project area.

**Table 2. Yakus EAA (3,987 acres)**

<b>Time Frame</b>	<b>Road Density (mi/mi<sup>2</sup>)</b>	<b>Security (%)</b>	<b>Openings (acres)</b>	<b>EHE (%)</b>
Existing	2.3	15	216	40
Proposed Action	2.5	7	617	38
Post Action	2.4	13	617	39

The Yakus EAA would gain 401 acres of potential forage for elk from harvest treatments in the proposed action. During implementation of the proposed action, elk security would decrease due to the loss of hiding cover in harvested units, and the use of some previously closed roads and construction of temporary roads to access harvest units. Elk would move away from the disturbance created by timber harvest or burning activities to other security areas. Elk are known to return to harvest units during hours of darkness to forage on vegetation. After implementation, roads would return to the existing condition status: (open or closed to motorized access).

A livestock allotment occupies a small presence (one square mile) in the EAA. Twenty head of cattle may use this area for about one-third of the year. The potential competition from cattle on forage, drops elk use of this area by 4%. This is a constant for the EAA and is carried through all time frames of analysis that include the existing condition, during implementation, and post project activities.

Elk vulnerability is linked to hunter and motorized route densities to predict elk mortality rates. The scale used for evaluating elk vulnerability is at the Game Management Unit (GMU (The Yakus EAA contributes to 0.4% of acres for the Dworshak GMU.) This GMU contains over 995,000 acres and includes mixed ownership, high road densities, and high bull elk vulnerability. The GMU is managed for high levels of hunter opportunity and moderate bull elk quality as accepted by past hunters in the zone.

This project would not increase road or hunter densities across the 1,556 square miles in the Dworshak Zone, and no measurable change would occur for elk mortality of survival. Therefore this project would not affect elk vulnerability in the GMU.

The temporary and other roads opened for harvest activities would not be maintained for public motorized access, and such use is discouraged. However, hunters may access the area by non-motorized means to hunt big game. The openings created by timber harvest may provide hunting opportunities for elk.

Upon completion of timber harvest, planting would occur to restock the units, and temporary roads would be decommissioned. Vegetative screening would increase (10-15 years) to a density where an elk would not be visible to a human that could be a distance of 200 feet or greater from the animal. As this "hiding" cover increases, it adds to the amount of elk security in the area. Increased elk security would also occur from the decommissioning of temporary roads. Together, these factors reduce elk vulnerability.

All temporary roads used during the implementation of the project would be decommissioned upon completion of project activities. Elk vulnerability would decrease to the same levels as the existing condition as vegetative hiding cover recovers.

## **Cumulative Effects**

Activities proposed for the project area may disturb or temporarily displace individual elk. The cattle presence in the Yakus EAA would not change, and elk use of the small portion of the livestock allotment would remain at 96%. Few roads are open to the public motorized traffic: about 15 miles in the Yakus EAA. The proposed action would increase openings (future forage habitat) by 10% in the EAA. No cumulative effects from other projects would occur in the EAA.

Firewood cutting would continue along the few roads in the project area. This activity would have an immeasurable effect, as only dead or dying trees along a road open to motorized traffic may be affected but could reduce security for elk. Fire suppression would continue, which would reduce the recruitment of early seral habitat from an uncontained wildfire. Both activities may disturb or displace elk.

## **Conclusion**

Some disturbance to elk would occur during the implementation of the action alternative. The Stray Creek Project would increase forage habitat for elk, while meeting the recommended guidelines for managing elk habitat. Project design criteria (WL-1) would minimize disturbance during the elk calving season. No measurable change in Elk Vulnerability from the proposed project was evident. The Yakus EAA would continue to provide EHE levels well above the 25% standard.

## ***Shiras Moose***

Moose are a management indicator species for big game species and old-growth/Pacific yew habitats. Potential habitat (Pierce and Peek, 1984) was queried as grand fir at or above 15 inches dbh. Current potential moose habitat in the project area is about 270 acres.

## **Direct and Indirect Effects**

The proposed action would reduce potential moose habitat in the project area by about 134 acres. Activities that would reduce tree habitat include temporary roads, timber harvest and burns in units that would be replanted with favored tree species. In 3-15 years, forage for moose would be available as understory herbs and shrubs recover. Hiding cover would be available after 15 years and would offer screening of an individual moose at 200 feet or greater from an open road. Project activities may disturb or displace an individual moose. Operations would occur during daylight hours, and moose may wander back into the units to feed on vegetation that has become available from logging operations during periods of darkness. About 85 acres of additional habitat is adjacent to the project area. This would offer potential displacement habitat for a moose.

## **Cumulative Effects**

The cumulative effects analysis area is the project area. Timeframe for recovery of potential moose habitat from a timber harvest is about 50 years: the period for a tree to attain the size and cover that contributes to moose habitat. Wildfire suppression would continue and may reduce the size of moose habitat lost if the fire was not contained.

## **Conclusion**

The proposed action would reduce potential habitat for up to approximately 15 years until hiding cover is established again; project activities may disturb and displace an individual moose.

## References

- Aney, W. C., & McClelland, B. R. (1990). Old-growth habitats and associated wildlife species in the northern Rocky Mountains (R1-90-42). Unpub. Report, USDA, Forest Service, Missoula MT.
- Arnett, E.B. & J.P. Hayes. 2009. Use of conifer snags as roosts by female bats in Western Oregon. *J. of Wildl. Mgmt.* 73(2):214-225.
- Badry, M. J. (2004). Fisher (*Martes pennanti*). Accounts and Measures for Managing Identified Wildlife—Accounts V. B.C. Min. of Water, Land and Air Protection. 19pp.
- Barclay, R.M.R. & A. Kurta. 2007. Ecology and behavior of bats roosting in tree cavities and under bark. in *Bats in Forests, Conservation and Management*. Eds. Lacki, M.J., J.P. Hayes & A. Kurta. pp. 17-59.
- Bartelt, P.E., C.R. Peterson, and R.W. Klaver. 2004. Sexual difference in the post-breeding movements and habitats selected by western toads (*Bufo boreas*) in southeastern Idaho. *Herpetologica* 60(4): 455-467.
- Bonar, L. 2001. Pileated woodpecker habitat ecology in the Alberta foothills. Thesis, Univ. of Alberta, Edmonton, Alberta, Canada.
- Boyce, M. S., Mao, J. S., Merrill, E. H., Fortin, D., Turner, M. G., Fryxell, J., & Turchin, P. (2003). Scale and heterogeneity in habitat selection by elk in Yellowstone National Park. *Ecoscience*, 10(4), 421-431.
- Brewer, L.T., R. Bush, J.E. Canfield, & A.R. Dohmen. 2009. Northern goshawk northern region overview. USDAFS, Region 1, Missoula, MT. 54pp.
- Brown, S. R. (2015). Nez Perce-Clearwater National Forests VMap 2014. USDA Forest Service, Region 1. 9pp.
- Brown, S., & Barber, J. (2012). The Region 1 Existing Vegetation Mapping Program (VMAP) Flathead National Forest Overview (p. 5). Version 12. Rep. US Forest Service Region 1, 12-34.
- Bull, E. L., Nielsen-Pincus, N., Wales, B. C., & Hayes, J. L. (2007). The influence of disturbance events on pileated woodpeckers in Northeastern Oregon. *Forest ecology and management*, 243(2-3), 320-329.
- Bull, E. L. 2006. Sexual differences in the ecology and habitat selection of western toads (*Bufo boreas*) in northeastern Oregon. *Pacific Northwest Research Station La Grande, OR. Herpetological Conservation and Biology* 1(1):27-38.
- Bull, E. L., Heater, T. W., & Shepherd, J. F. (2005). Habitat selection by the American marten in northeastern Oregon. *Northwest Science*. 79 (1): 36-42.
- Bull, E. L., and J. A. Jackson. 1995. Pileated woodpecker (*Dryocopus pileatus*). No. 148. A. Poole, and F. Gill, editors. *The Academy of Natural Sciences*, Philadelphia, Pennsylvania, and *The American Ornithologists' Union*, Washington, DC, USA.
- Bull, E. L., and R. S. Holthausen. 1993. Habitat use and management of pileated woodpeckers in northeastern Oregon. *Journal of Wildlife Management* 57: 335-345.

- Bull, E.L., S. Holthausen & M.G. Henjum. 1992. Roost trees used by pileated woodpeckers in Northeastern Oregon. *J. Wildl. Mgmt* V. 56(4):786-793.
- Busing, R. T., Halpern, C. B., & Spies, T. A. (1995). Ecology of Pacific yew (*Taxus brevifolia*) in western Oregon and Washington. *Conservation Biology*, 9(5), 1199-1207.
- Buskirk, S. W., & Ruggiero, L. F. (1994). American marten. In L. F. Ruggiero, K. B. Aubry, S. W. Buskirk, L. J. Lyon, & W. J. Zielinski (Eds.), *The scientific basis for conserving forest carnivores: American marten, fisher, lynx, and wolverine* (Gen. Tech. Rep. RM-254) (pp. 7-37). Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station.
- Chapin, T., G., Harrison, D. J., & Katnik, D. D. (1998). Influence of landscape pattern on habitat use by American marten in an industrial forest. *Conservation Biology*, 12(6), 1327-1337.
- Christensen, A.G., L.J. Lyon & J.W. Unsworth. 1993. Elk management in the Northern Region: considerations in forest plan updates or revisions. USDAFS Intermountain Res. Sta. GTR INT-303.
- Cook, J.G., L.L. Irwin, L.D. Bryant, R.A. Riggs & J.W. Thomas. 1998. Relations of forest cover and condition of elk: a test of thermal cover hypothesis in summer and winter. *Wildl. Monographs*, n 141:3-61.
- Federal Register. 2001. Executive Order 13186. Responsibilities of federal agencies to protect migratory birds. FR(v)66, No. 11. 1-17-2001. 3853-3856.
- Frair, J.L., E.H. Merrill, H.L. Beyer & J.M. Morales. 2008. Thresholds in landscape connectivity and mortality risks in response to growing road networks. *J. of Appl. Ecol.* 45:1504-1513.
- Giese, C.L., & F.J. Cuthbert, 2003. Influence of surrounding vegetation on woodpecker nest tree selection in oak forests of the Upper Midwest, USA. *For. Ecology & Mgmt*, 179(1), 523-534.
- Guscio, C. G., B.R. Hossack, L.A. Eby, and P.S. Corn. 2007. Post-breeding habitat use by adult boreal toads (*Bufo Boreas*) after wildfire in Glacier National Park, USA. *Herp. Cons. Biol.* 3 (1): 55-62.
- Hargis, C. D., Bissonette, J. A., & Turner, D. L. (1999). The influence of forest fragmentation and landscape pattern on American martens. *Journal of Applied Ecology*, 36(1), 157-172.
- Hutto, R.L. 1995. Composition of bird communities following stand-replacement fires in northern Rock Mountain (USA) conifer forests. *Cons. Biol.* V.9(5):1041-1058.
- Idaho Department of Fish and Game. 2017. Wolf Statewide Report. 21 pp. Boise, ID
- Idaho Dept. of Fish and Game. 2016. Moose Report. 78 pp. Boise, ID.
- Idaho Department of Fish and Game and Nez Perce Tribe. 2014. 2013 Idaho wolf monitoring progress report. Idaho Department of Fish and Game, 600 South Walnut, Boise, Idaho; Nez Perce Tribe Wolf Recovery Project, P.O. Box 365, Lapwai, Idaho. 74 pp.
- Johnson, J. S., M. J. Lacki, and M.I D. Baker. 2007. "Foraging ecology of long-legged myotis (*Myotis volans*) in north-central Idaho." *Journal of Mammalogy* 88, no. 5 (2007): 1261-1270.

- Kennedy, P.L. (2003, January 2). Northern Goshawk (*Accipiter gentiles atricapillus*): a technical conservation assessment. [Online]. USDA Forest Service, Rocky Mountain Region. Available: <http://www.fs.fed.us/r2/projects/scp/assessments/northerngoshawk.pdf>
- Kirk, D. A., & Naylor, B. J. (1996). Habitat requirements of the Pileated Woodpecker (*Dryocopus pileatus*) with special reference to Ontario. Ontario Ministry of Natural Resources (No. 46, p. 49). SCST Technical Report.
- Koehler, G. M., & Hornocker, M. G. (1977). Fire effects on marten habitat in the Selway-Bitterroot Wilderness. *The Journal of Wildlife Management*, 500-505.
- Koehler, G. M., Moore, W. R., & Taylor, A. R. (1975). Preserving the pine marten: management guidelines for western forests. *Western Wildlands*.
- Lacki, M. J., Baker, M. D., & Johnson, J. S. (2010). Geographic variation in roost- site selection of long- legged myotis in the Pacific Northwest. *The Journal of Wildlife Management*, 74(6), 1218-1228.
- Lacki, M. J., Johnson, J. S., Dodd, L. E., & Baker, M. D. (2007). Prey consumption of insectivorous bats in coniferous forests of north-central Idaho. *Northwest Science*, 81(3), 199-205.
- Montgomery, R.A., G.J. Roloff & J.J. Millspaugh. 2012. Variation in elk response to roads by season, sex and road type. *J. of Wildl. Mgmt.* 77(2):313-325.
- Moriarty, K. M., Zielinski, W. J., & Forsman, E. D. (2011). Decline in American marten occupancy rates at Sagehen Experimental Forest, California. *The Journal of Wildlife Management*, 75(8), 1774- 1787.
- Moser, B. W., & Garton, E. O. (2019). Northern goshawk space use and resource selection. *The Journal of Wildlife Management*, 83(3), 705-713.
- Moser, B. W., & Garton, E. O. (2009). Short-term effects of timber harvest and weather on Northern Goshawk reproduction in northern Idaho. *Journal of Raptor Research*, 43(1), 1-11.
- Moser, B.W. 2007. Space use and ecology of goshawks in Northern Idaho. Dissertation in Natural Resources, U of I, Moscow, Idaho.
- Potvin, F., Bélanger, L., & Lowell, K. (2000). Marten habitat selection in a clearcut boreal landscape. *Conservation Biology*, 14(3), 844-857.
- Nadeau, M. S., DeCesare, N. J., Brimeyer, D. G., Bergman, E. J., Harris, R. B., Hersey, K. R., ... & Thomas, T. P. (2017). Status and trends of moose populations and hunting opportunity in the western United States. *Alces: A Journal Devoted to the Biology and Management of Moose*, 53, 99-112.
- NatureServe. 2019. Online database that tracks species status primarily in North America.
- Olson, L. E., Sauder, J. D., Albrecht, N. M., Vinkey, R. S., Cushman, S. A., & Schwartz, M. K. (2014). Modeling the effects of dispersal and patch size on predicted fisher (*Pekania [Martes] pennanti*) distribution in the US Rocky Mountains. *Biological Conservation*, 169, 89-98.
- Pierce, D. J., & Peek, J. M. (1984). Moose habitat use and selection patterns in north-central Idaho. *The Journal of wildlife management*, 1335-1343.

- Poole, K. G., & Stuart-Smith, K. (2005). Finescale winter habitat selection by moose in interior montane forests. *Alces*, 41(1), 1-8.
- Raley, C.M., E.C. Lofroth, R.L. Truex, J.S. Yaeger, & J.M. Higley. 2012. Habitat ecology of fishers in Western North America. In *Biology and Conservation of martens, sables and fishers*. Editors: Aubry, Zielinski, Raphael, Proulx & Buskirk. Chapter 10: 231-254.
- Ranglack, D.H, K.M. Proffitt, J. Gude, J. Canfield, J. Rotella, & R.A. Garrott. 2017. Security areas for elk during archery and rifle hunting seasons. *J. Wildl Mgmt.* v.81(5):778-791.
- Rich, T.D., 2007. Guide to the Partners in Flight Population Estimates Database. Version: North American Landbird Conservation Plan 2004. Partners in Flight Technical Series No 5. <http://www.partnersinflight.org/>
- Rowland, M.M., M.J. Wisdom, B.K. Johnson & M.A. Penninger. 2005. Effect of roads on elk: implications for management in forested ecosystems. Pages 42-52 in Wisdom, M. J., technical editor, *The Starkey Project: a synthesis of long-term studies of elk and mule deer*.
- Rowland, M.M., M.J. Wisdom, B.K. Johnson & J.G. Kie. 2000. Elk distribution and modeling in relation to roads. *J. of Wildl. Mgmt.* v.64(3):672-684.
- Ruggiero, L.F., K.B. Aubry, S.W. Buskirk, L.J. Lyon and W.J. Zielinski (tech. eds.). 1994. The Scientific basis for conserving forest carnivores: American marten, fisher, lynx and wolverine in the western United States. General Technical Report RM-GTR-254. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 184 pp.
- Sauder, J.D. 2014. Landscape ecology of fishers (*Pekania pennant*) in North-Central Idaho. Dissertation in Natural Resources. U of Idaho, Moscow, Idaho.
- Sauder, J.D. and J.L. Rachlow. 2015. Forest heterogeneity influences habitat selection by fishers (*Pekania pennant*) within home ranges. *For. Ecol. & Mgmt.* 347: 49-56.
- Sauder, J.D. and J.L. Rachlow. 2014. Both forest composition and configuration influence landscape-scale habitat selection by fishers (*Pekania pennanti*) in mixed coniferous forests of the Northern Rocky Mountains. *Forest Ecology and Mgmt.* 314: 75-84.
- Schultz, C. 2010. Effective wildlife conservation planning. *BioScience.*, V. 60, N.7, 545-551.
- Schwartz, M. K., DeCesare, N. J., Jimenez, B. S., Copeland, J. P., & Melquist, W. E. (2013). Stand-and landscape-scale selection of large trees by fishers in the Rocky Mountains of Montana and Idaho. *Forest Ecology and Management*, 305, 103-111.
- Servheen, G. 1997. Interagency guidelines for evaluating and managing elk habitats and populations in Central Idaho. 67+pp.
- Shirk, A. J., Raphael, M. G., & Cushman, S. A. (2014). Spatiotemporal variation in resource selection: insights from the American marten (*Martes americana*). *Ecological Applications*, 24(6), 1434-1444.
- Squires, J. R., and P. L. Kennedy. 2006. Northern goshawk ecology: an assessment of current knowledge and information needs for conservation management. *Studies in Avian Biology* 31: 8-62.
- Squires, J.R. & R.T. Reynolds. 1997. Northern goshawk. *Birds of North America*, No. 298. Pp. 31.

- Thomas, J.W., H. Black, R.J. Scherzinger & R.J. Pedersen. 1979. Deer and Elk, in Wildlife habitats in managed forests, the Blue Mountains of Oregon and Washington. 104-127.
- Unsworth, J.W., L. Kuck, E.O. Garton, and B.R. Butterfield. 1998. Elk habitat selection on the Clearwater National Forest, Idaho. *Journal of Wildlife Management*. 62(4): 1255-1263.
- Unsworth, J.W., L. Kuck, M.D. Scott, & E.O. Garton. 1993. Elk mortality in the Clearwater drainage of Northcentral Idaho. *J. Wildl. Mgmt* 57(3):495-502.
- USDA Forest Service. 2008. Memorandum of understanding between the U.S. Department of Agriculture Forest Service and the U.S. Fish and Wildlife Service to promote the conservation of migratory birds. FS Agreement #08-MU-1113-2400-264. 13pp.
- USDA Forest Service. 2019. Nez Perce-Clearwater National Forests Plan Revision Draft Environmental Impact Statement. Chapter 3, 3.2.3.2 Fisher pp. 83-88.
- USDI Fish and Wildlife Service. 1998. Northern Goshawk Status Review. US Fish and Wildlife Service, Portland, Oregon, USA
- USDI Fish and Wildlife Service. 2017. Species Status Assessment for the Northern Rocky Mountains Fisher (V.2). U.S. Fish and Wildlife Service, Denver, Co. 83pp.
- USDI Fish and Wildlife Service. 1987. Northern Rocky Mountain Wolf Recovery Plan. Denver, CO. 119pp.
- Vonhof and Barclay. 1997. Use of tree stumps as roosts by the western long-eared bat. *J. Wildl. Mgmt*. 61(3):674-684.
- Wasserman, T. N., Cushman, S. A., Schwartz, M. K., & Wallin, D. O. (2010). Spatial scaling and multi-model inference in landscape genetics: *Martes americana* in northern Idaho. *Landscape ecology*, 25(10), 1601-1612.
- Wasserman, T. N., Cushman, S. A., Shirk, A. S., Landguth, E. L., & Littell, J. S. (2012). Simulating the effects of climate change on population connectivity of American marten (*Martes americana*) in the northern Rocky Mountains, USA. *Landscape Ecology*, 27(2), 211-225.
- Weir, R.D. & A.S. Harestad. 2003. Scale dependent habitat selectivity by fishers in South-Central British Columbia. *J. of Wildl. Mgmt*. 67(1):73-82.
- Weir, R. D., & Corbould, F. B. (2010). Factors affecting landscape occupancy by fishers in north-central British Columbia. *Journal of Wildlife Management*, 74(3), 405-410.
- Wisdom, M.J, R.S. Holthausen, B.C. Wales, C.D. Hargis, V.A. Saab, D.C. Lee, W.J. Hahn, T.D. Rich, M.M. Rowland, W.J. Murphy, & M.R. Eames. 2000. Source habitats for terrestrial vertebrates, of focus in the interior Columbia Basin: broad-scale trends and management implications. Volume 1.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [http://www.ascr.usda.gov/complaint\\_filing\\_cust.html](http://www.ascr.usda.gov/complaint_filing_cust.html) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

USDA is an equal opportunity provider, employer and lender.